

EFFECTS OF DIETARY SUPPLEMENTATION OF *Prosopis africana* (African mesquite) ESSENTIAL OIL ON THE GROWTH PERFORMANCE, NUTRIENT RETENTION AND OXIDATIVE STRESS INDICES OF BROILER CHICKEN

Alagbe, John Olujimi^{1*}, Agubosi, Oluchi Precious² & Oluwafemi, Rufus Adebisi

¹Department of Animal Science, University of Abuja, Gwagwalada, Nigeria

Corresponding Author: Alagbe, Olujimi John E-mail: dralagbe@outlook.com

ARTICLE INFO

Received: May 25, 2023

Accepted: August 4, 2023

Volume: 3

Issue: 3

KEYWORDS

Broilers, Phytochemicals, Performance, Nutrient, Oxidative stress, *Prosopis africana*

ABSTRACT

This research was undertaken to evaluate the effects of dietary supplementation of *Prosopis africana* essential oil on the growth performance, nutrient retention and oxidative stress indices of broiler chickens. A total of 540 one-day-old (Arbo acre) broiler chicks were randomly assigned to six treatment groups with six replicates of fifteen birds each. Experimental diet was adequate in all nutrients according to NRC (1994) recommendation. Dietary treatments were as follows: treatment 1 (T₁): basal diet with no *Prosopis africana* oil (PAOs); T₂: basal diet plus neomycin at 2.5 g/kg while T₃, T₄, T₅ and T₆ were fed basal diet supplemented with 200 mg, 400 mg, 600 mg and 800 mg/kg PAOs respectively. Improvements for body weight gain and feed conversion ratio ($P < 0.05$) were observed in birds fed PAOs compared to the other treatments. Mortality rate was highest in T₁ (5.33 %) followed by T₂ (3.02 %) and T₃ (1.00 %) ($P < 0.05$). Dressing percentage and weights of organs were significantly ($P < 0.05$) influenced by the treatments except the weights of kidney, spleen, pancreas and bursa ($P > 0.05$). Dry matter and crude protein digestibility were highest in T₃ through T₆, intermediate in T₂ and lowest in T₁ ($P < 0.05$). Malondialdehyde, catalase, glutathione peroxidase, reduced glutathione and superoxide dismutase were significantly ($P < 0.05$) different among the treatments. It was concluded that the dietary supplementation of PAOs up to 800 mg/kg in broiler chickens could arouse its appetite translating to higher weight gain, increased secretion of enzymes in the gastro intestinal tract as well as reduction of free radicals.

1. Introduction

Phytogenics are substances that come from plants such as spices, herbs, essential oils and other plant extracts. They have been utilized for countless years for both traditional uses and for their unique abilities as natural treatments (Tobias, 2010). Among other biological actions, phytogenic substances have antioxidant, antibacterial, antifungal, antiviral, hepato-protective, immune-stimulatory amongst others (Alagbe, 2022; Singh *et al.*, 2021). Following the 2009 European Union ban on antimicrobial growth promoters, numerous studies were conducted to investigate possible replacements. Using essential oils have been adopted as one of the potential alternatives to antibiotics (Musa *et al.*, 2020; Axel, 2018).

Essential oils are volatile oils extracted from plants using steam or hydrodistillation as the distillation method (Marc, 2010). They contain plant – derived chemicals with a sophisticated chemical make-up from leaves, stems, bark, seeds, flowers and other plant parts essential oils can be extracted (Alagbe *et al.*, 2023; Axel, 2018). It is thought that they hold knowledge and specific instructions for the conduct and survival of the plants (Tobias, 2010; Benzoni and Guyonvarch, 2012). Additionally, the harvest season, climatic conditions and cultivar within a plant variety – particularly for *Prosopis africana* oil have an impact on the composition of essential oils (Alagbe, 2022; Sandra, 2018).

The essential oil of *Prosopis africana* is packed with volatile and fragrant substances. Prosogerin is one of the oil's major bioactive components which has been discovered to play a part in processes such as stimulation of digestion, antimicrobial properties, antioxidant activity and stimulation of feed intake (Sandra, 2018; Jan *et al.*, 2013). In broilers, *in vivo* antimicrobial efficacy of *Prosopis africana* oil on *Escherichia coli*, *Staphylococcus spp*, *Streptococcus spp* and other pathogenic organisms when supplemented at 300 mg/kg was demonstrated (Shittu and Alagbe, 2010). Furthermore, in an experiment carried out by Alagbe *et al.* (2022), dietary inclusion of *Prosopis africana* oil (PROs) at 400 mg/kg for broilers stimulate digestive process, efficient feed utilization and balanced gut microflora. Field trials have shown the great benefit of PROs application in attenuating inflammatory processes in the intestine by blocking the pro-inflammatory transcription factor NF-kB and removing free radicals in birds when included in the diet at 100 mg/kg (Rao, 2015). It has been reported that essential oils can act as immunosuppressors or immunostimulators to stimulate the immune system through the host (Tobias, 2010; Alagbe, 2017). They can also the levels of plasma lipid peroxidation and malondialdehyde, it can help to lessen excessive oxidative stress (Marc, 2010; Luis, 2012). Reactive oxygen species play a role in a variety of diseases, including chronic inflammation, hence there is a close relationship between antioxidant and anti-inflammatory function (Adewale *et al.*, 2021).

Prosopis africana oil has a lot of promise and it's important to set up a foundation for adding it to broiler's diet. The purpose of this experiment was to ascertain how well broiler chickens given *Prosopis africana* essential oil performed in terms of growth, carcass characteristics, nutrient digestibility and oxidative stress markers.

2. Methodology

2.1 Location of the experiment, ethical approval and *Prosopis africana* essential oil extraction

The experiment was conducted at the University of Abuja Teaching and Research Farm, Department of Animal Science along Airport Road in Gwagwalada which is positioned between latitudes 8°57¹ and 8°55¹N and longitudes 7°05¹ and 7°06¹E (NPC,

2006). The department of Animal Science at the University of Abuja's research ethics committee accepted the protocol for the study, which was conducted in accordance with its specifications and rules.

The *Prosopis africana* seeds utilized in this study was purchased from a local market in Gwagwalada in April, 2021. With a voucher number of ANS/203ER/008, the seeds were identified and verified at the Department of Biological Sciences herbarium, University of Abuja, Nigeria. *Prosopis africana* oil was obtained using the steam-distillation method, which also calls for the use of a digital scale, a round-bottom flask, a heating mantle made of glass fiber, a measuring cylinder, and a separatory funnel. 150 g of ground *Prosopis* seed and 250 mL of distilled water were put into a flask with a circular bottom. The resulting mixture was transferred to a glass yarn heating mantle and heated to an internal temperature of 80 degree celsius before the condenser was placed on top of the flask with a circular bottom. Fifteen minutes of forced boiling is given to the combination, after which the distillate is collected in a beaker until no more oil drops are present. The distillate was introduced to the separatory funnel in order to obtain *Prosopis africana* essential oil.

2.2 Animal care (diets, housing, and arrangements)

540 mixed-sex 1-day-old broiler chicks (Arbo Acres), obtained from a well-known commercial hatchery in Ogun State, Nigeria, were divided into six treatments, each with six replicates of fifteen birds. Galvanized hand feeders and automatic nipple drinkers were installed in a galvanized battery cage that measured (100 cm × 75 cm × 50 cm) (length depth height) in semi-open pens. Chicks were given anti-stress (glucose plus water) at a rate of 2 grams of glucose to 10 liters of water after being weighed upon arrival to ascertain their average starting body weight. In this investigation, an entirely random experimental design was adopted. Diets were compounded based on NRC (1994) nutritional needs for broilers, which are shown in Table 2. The experimental diet provides 23.30% crude protein (CP), 21.40% CP and metabolizable energy content (ME) of 2991.5 Kcal/kg and 3108.1 Kcal/kg respectively at the starter phase (1-28 days) and finisher phase (29-56 days). For the duration of the experiment's eight-week run, all birds had free access to food and water.

The following is a representation of the experimental arrangement:

Treatment one (T1): Basal diet with no *Prosopis africana* oil (PAOs)

Treatment two (T2): Basal diet plus 2.5 g/kg of Neomycin

Treatment three (T3): Basal diet plus 200 mg/kg *Prosopis africana* oil

Treatment four (T4): Basal diet plus 400 mg/kg *Prosopis africana* oil

Treatment five (T5): Basal diet plus 600 mg/kg *Prosopis africana* oil

Treatment six (T6): Basal diet plus 800 mg/kg *Prosopis africana* oil

2.3 Variables collected

2.3.1 Performance traits

The difference between the feed offered and rejected was used to calculate the amount of feed consumed. The ratio of average feed intake to average body weight growth was used to figure out the feed conversion ratio, or the amount of feed required to create one unit of gain. The difference between the end body weight and the starting body weight was used to compute weight gain. Weight gain for each treatment divided by the number of experimental days was used to calculate the average daily body weight.

The number of deceased birds divided by the total number of birds multiply by 100 was used to compute the mortality percentages all through the experimental period.

2.3.2 Carcass quality evaluation

12 birds from each treatment were randomly chosen at the close of the experiment to be evaluated for carcass quality. In order to avoid contamination and fly invasion, carcass examination was done in a clean, fly-free environment. Prior to slaughter, the chosen birds were weighted and feed-starved for a twelve-hour period. By cutting the jugular vein, putting the bird into hot water, and manually plucking the feathers, the birds were humanely killed. The heart, pancreas, kidney, lungs, and proventriculus were carefully removed from the carcass. Each organ was then weighed separately and documented.

The ratio of carcass weight to slaughter weight, multiplied by 100, was used to compute the dressing %. The eviscerated carcass's breast, wings, thighs with drumsticks, and backside were all removed, weighed, and the percentage of the remaining weights of each were calculated by dividing the total weight by the slaughter weight and multiplying the result by 100.

2.3.3 Nutrient retention trial

At the conclusion of the study (day 56), twelve birds were randomly chosen from the treatment and placed in individual metabolic cages (measuring 25 cm × 20 cm × 30 cm in length, depth, and height) with automatic nipple drinkers and galvanized manual feeders. Birds were acclimated and fed a known amount of food on the first day of the trial. Droppings were collected, weighed, and air dried daily for each replicate. Faecal excretions were bulked up for each replicate on the final day of the study and delivered to the lab for additional analysis. Proximate analysis of faecal droppings were carried out using near infra-red spectroscopy (NIRS™ DA1600) with specifications: dimension (200 × 130 × 100 mm) (w×d×h), optical band width (9.44± 0.76 nm), wave length (1400 - 2000 nm), spectral resolution (0.9 nm /data point), detector (206 pixel in GaAs diode array), absorbance range (up to 3 AU) and wavelength precision (< 0.08 nm). The data were expressed as follows:

Nutrient retention (% DM) is equal to the difference between intake and output in the form of excretion divided by intake multiplied by 100.

2.3.4 Oxidative stress investigation

Samples of liver were harvested from 12 birds per treatment (from birds used for carcass investigation). Liver samples were immersed into potassium solution and subjected to test using NEXRO kit (Model 81-09A, India). The kit has the following components; (DNPH solution, hematoxylin, wash buffer, biotinylated secondary antibody, antigen retrieval buffer). Oxidative detection of glutathione peroxidase, catalase, reduced glutathione, superoxide dismutase and melondialdehyde were measured on the display or output unit.

2.4 Statistical investigation

Using Statistical Package for Social Sciences [SPSS (18.0)], one-way analysis of variance was performed on all data, and Duncan multiple range tests were used to identify significant means. If $P < 0.05$, significance was deemed to exist.

Table 1: Gross composition of experimental diets

Components	Starters phase (0-28 days)	Finishers phase (29-56 days)
Maize	53.50	59.00
Wheat offal	2.00	5.00
Soya bean meal	30.00	24.00
Groundnut cake	7.00	5.00
Fish meal (65 %)	2.00	1.00
Limestone	1.50	2.00
Bone meal	3.00	4.00
Lysine	0.20	0.20
Methionine	0.20	0.20
*Premix	0.25	0.25
Salt	0.30	0.35
***Toxin binder	0.10	0.10
Total	100.0	100.0
Determined analysis		
Crude protein (%)	23.30	21.40
Crude fibre (%)	4.18	5.01
Ether extract (%)	4.03	4.47
Calcium (%)	1.50	1.80
Phosphorus (%)	0.58	0.66

Energy (kcal/Kg)	2991.5	3108.1
------------------	--------	--------

*Premix supplied per kg diet: - vit A, 14,000 I.U; vit E, 5mg; vit D3, 3000I.U, vit K, 3mg; vit B2, 5.5mg; Niacin, 25mg; vit B12, 16mg; choline chloride, 120mg; Mn, 5.2mg; Zn, 25mg; Cu, 2.6g; folic acid, 2mg; Fe, 5g; pantothenic acid, 10mg; biotin, 30.5g; antioxidant, 56mg (starter's mash)

**Premix supplied per kg diet: - vit A, 9,000 I.U; vit E, 10mg; vit D3, 1500I.U, vit K, 3.8mg; vit B2, 10 mg; Niacin, 15mg; vit B12, 10mg; choline chloride, 250mg; Mn, 5.0mg; Zn, 56mg; Cu, 1.6g; folic acid, 2.8mg; Fe, 5.1g; pantothenic acid, 10mg; biotin, 30.5g; antioxidant, 56mg (finisher's mash)

*** Toxin binder composition per Kilogram: Mannan Oligo Saccharide (MOS) (50gm), Activated charcoal 20 grams, activated charcoal 20 grms, ammonium format 20 grams, citric acid 10 grams, calcium propionate 20 grams, sodium bentonite 20 grams, hydrated sodium calcium aluminosilicate 725 grams.

3.0 Results and Discussion

3.1 Performance of broiler chickens fed diet containing *Prosopis africana* essential oil (PAOs)

Growth performance of broiler chickens fed *Prosopis africana* oil (Table 2) revealed that the initial body weight, final body weight, body weight gain, average daily weight gain, total feed intake, average daily feed intake, feed conversion ratio and mortality varied between 43.29 - 44.20 grams, 1825.1 - 3024.6 grams, 1029.6 - 1775.6 grams, 18.39 - 31.71 grams, 4307.2 - 4425.6 grams, 76.91 - 79.03 grams, 1.46 - 2.40 and 1.00 - 8.33 percent respectively. The treatments considerably differed ($P<0.05$) in average daily weight gain (ADWW), feed conversion ratio and % mortality. Treatment 4 (T4) through T6 have the highest ADWW followed by T3 and lowest in T1 and T2 ($P<0.05$). Conversely, mortality percentage were maximum in T1 compared to the other groups ($P<0.05$). Average daily feed intake values were not statically influenced ($P>0.05$) by the treatments, though it increased slightly through treatment 1 to 6 but not at a significant rate. The experimental diets were formulated to meet the nutrient requirements of broiler chicks as recommended by NRC (1994). In this study, the presence of phytochemicals or secondary metabolites in *Prosopis africana* oil could cause beneficial effects on their performance and health status of broiler chickens. In this study, the improved body weight gain (ADW) and feed conversion ratio (FCR) of the experimental birds fed *Prosopis africana* oil supplemented diets could be due to the activities of the constituents of these phytogenics (Oluwafemi *et al.*, 2021; Muritala *et al.*, 2022). Among the treatments, birds fed higher doses of PAOs (300 and 400 mg/kg) gained more weight than those on lower dose (100 and 200 mg/kg), implying that higher dose enhanced feed utilization better than the other treatments. The multiple biological activities of phytochemicals could have contributed to the improved BWg and FCR recorded in this group of birds (Alagbe *et al.*, 2023; Singh *et al.*, 2021). Similar findings were made by Hafeez *et al.* (2016), who showed that matrix encapsulation of essential oils with active ingredients like thymol and limonene at the dosage of 100 mg/kg feed improved performance and apparent ileal digestibility of nutrients in broiler chickens compared with control. In addition, broiler chicks fed 0.5 mg/kg of anise seed oil with anethole as the active ingredient significantly increased body weight gain while having no impact on feed intake or feed conversion ratio, according to Soltan *et al.* (2008). Furthermore,

broiler chicks fed a higher level of anise seed oil (1.5 mg/kg) had lower growth performance in comparison to controls. This range of findings may indicate that phytogetic feed additives differ significantly in their chemical makeup depending on their constituents, climatic conditions, location, age of plants, dosage used, processing techniques, and conditions during harvest or storage (Agubosi *et al.*, 2021). The lowest weight increase was shown in the T1 group of birds, which did not receive any PAOs or antibiotics. This clearly shows that broilers cannot be raised in the tropics without the use of growth promoters, whether they be organic or synthetic.

According to reports, the PAOs increased the flavor and palatability of the feed, which led to an increase in voluntary feed intake and improved weight gain. This result contradicts Schone *et al.*'s (2006) findings, which showed that birds fed diets supplemented with caraway oil at 100 mg/kg consumed less feed overall. No mortality was observed in birds fed diets enriched with PAOs, suggesting that the oil may be able to stop the growth of pathogenic bacteria in the gastrointestinal tract. This outcome is consistent with the findings of Oloruntola *et al.* (2021), who supplemented the feed of broiler chicks with *Irvingia gabonensis* and *Ocimum gratissimum* powder.

Table 2: Performance of broiler chickens fed diet containing *Prosopis africana* essential oil

Treatments	IBw (g/bird)	FBw (g/bird)	BWg (g/bird)	ADWW (g/bird)	TFII (g)	ADFII (g/bird)	FCR	M (%)
T1	44.20	1825.1 ^d	1029.6 ^c	18.39 ^c	4307.2	76.91	2.40 ^a	5.33 ^a
T2	44.10	2464.7 ^c	1383.3 ^c	24.70 ^c	4419.0	78.91	1.80 ^b	3.02 ^b
T3	43.29	2744.1 ^b	1614.1 ^b	28.82 ^b	4403.1	78.63	1.60 ^{bc}	1.00 ^c
T4	44.08	2854.7 ^a	1708.6 ^a	30.51 ^a	4400.9	78.59	1.54 ^c	-
T5	44.10	2985.3 ^a	1736.9 ^a	31.02 ^a	4421.1	78.95	1.50 ^c	-
T6	43.93	3024.6 ^a	1775.6 ^a	31.71 ^a	4425.6	79.03	1.46 ^c	-
SEM	0.76	0.55	33.89	0.33	1.71	0.47	0.06	-
P-value	0.01	0.18	0.29	0.11	0.44	0.02	0.001	0.93

Values in sections with various characters differ considerably ($p < 0.05$); Initial Current Weight (IBw), Final Body weight (FBw), and gains in body weight (BWg) Average Daily Body Weight (ADWW), Average Daily Feed Consumption (ADFII), Feed Conversion Ratio (FCR), Mortality % (M), Standard Error of Mean (SEM), etc. Baseline diet + 0% *Prosopis africana* oil; Baseline diet + 2.0g/kg Neomycin; Baseline diet + 200 mg PAOs; Baseline diet + 400 mg PAOs; Baseline diet + 600 mg PAOs; Baseline diet + 800 mg PAOs.

3.2 Carcass characteristics of broiler chickens fed diets with PAOs

The impact of *Prosopis africana* oil on the carcass characteristics of broiler chicken is revealed in Tables 3. Final live weight (FLw), dress weight (Dw), dressing percentage (D %), and mean weight of the head, legs, gizzard, liver, lungs, heart, thigh, breast, back, kidney, spleen, bursa, and pancreas, respectively, varied from 1412.8 to 3200.4 grams, 967.8 to 2676.8 grams, 68.50 to 83.64% grams, 35.16 to 52.80 grams, 5.88 to 16.50 grams, 9.34 to 13.75 grams, 307.8 to 655.3 grams, 485.6 to 813.6 grams, 270.4 to 530.2 grams, 0.41 to 0.45 grams, 2.00 to 2.61 grams, 3.99 to 4.16 grams and 4.01 to 4.98 grams respectively. Treatment

3 through T6 had the highest dressing % values, whereas middle in T2 and T1 had the lowest value ($P<0.05$). In comparison with the remaining treatments, T4, T5 and T6 had the maximum weight for drumstick, legs, liver, lungs, heart, thigh, breast, and back ($P<0.05$). Treatments had no effect on the head, kidney, spleen, pancreas and bursa ($P>0.05$). The dressed weight usually expressed as dressing percentage is an indication of meat value that could be obtained from an animal. When compared to the other treatments, birds fed PAOs had the highest dressing % ($P<0.05$). This implies that adding *Prosopis africana* essential oil supplementation at a dose of 100 to 400 mg/kg could effectively substitute in-feed antibiotics without lowering the dressing %. Similar results were seen by Safa *et al.* (2014), who fed Hubbard broiler chickens garlic oil as a natural feed addition and observed enhanced carcass quality. The reduced organ weights, especially in the liver and kidney, show that the secondary metabolite concentrations of PAOs are not hazardous. According to Bamgbose *et al.* (2004), clothe weight and organ weight characteristics are actual symptoms or markers of the degree of anti-nutritional factor reduction. The absence of abnormal internal organ growth or atrophy suggests that the birds were able to tolerate PAOs. Since carcass features are a reflection of an animal's growth response, a decrease in dressed weight and dressing % is expected for (T1 without *Prosopis africana* oil or antibiotics) (Ose *et al.*, 2019; Adewale *et al.*, 2021).

Head, gizzard, kidney, pancreas, spleen, and bursa relative organ weights were not substantially ($P>0.05$) impacted by the treatments. As the supplemented amounts of PAOs rose across treatments, the relative weights of the lungs, heart, thigh, breast, drumstick, and back considerably ($P<0.05$) increased. The relative weight of gizzard, thigh and lungs has been reported to increase in birds supplemented with cinnamon oil at 500 mg/kg feed (Saied *et al.*, 2022), red pepper oil at 100 mg/kg (Reda *et al.*, 2020), Savory oil at 50 mg/kg (Sarica *et al.*, 2009) and oregano oil at 300 mg/kg (Krishan and Narang, 2014) while no changes were observed in birds fed rosemary oil (Attia *et al.*, 2017); rosemary extract (Al-hijazeen, 2021). According to Rougiere and Carre (2010) and Amerah *et al.* (2009), an increase in the relative weight of the gizzard in birds fed PAOs may lead to the swift production of pepsinogen and hydrochloric acid from the proventriculus, which will enhance feed degradation and reduce the entry of pathogenic bacteria into the digestive tract. Bile is produced by the liver, which is also connected to food metabolism (Amerah *et al.*, 2009; Alagbe *et al.*, 2019). The generation of digestive enzymes is carried out by the pancreas, whereas the spleen actively participates in filtering potentially hazardous chemicals before they enter the bloodstream (Van *et al.*, 1990).

Table 3: Effect of *Prosopis africana* oil on the retail cut parts and organ weights of broiler chicken

Parameters	T1	T2	T3	T4	T5	T6	SEM	P-value
FLW (g)	1412.8 ^d	1922.7 ^c	2500.8 ^b	2831.7 ^b	3100.4 ^a	3200.4 ^a	34.70	0.07
DW (g)	967.8 ^c	1481.1 ^b	2000.6 ^a	2291.8 ^a	2510.9 ^a	2676.8 ^a	20.81	0.04

DP (%)	68.50 ^c	77.03 ^b	80.00 ^a	81.01 ^a	81.00 ^a	83.64 ^a	5.02	0.22
Head (g)	41.22	43.05	45.60	45.11	46.02	46.07	1.18	0.03
D.stick (g)	123.7 ^c	166.1 ^b	178.6 ^{ab}	191.7 ^a	196.5 ^a	197.9 ^a	3.03	0.08
Leg (g)	50.28 ^c	78.44 ^{ab}	82.10 ^a	83.00 ^a	85.80 ^a	86.88 ^a	1.35	0.05
Gizzard (g)	68.67	69.05	70.94	71.60	71.49	71.04	1.20	0.25
Liver (g)	35.16 ^c	41.50 ^{ab}	50.47 ^a	52.39 ^a	52.01 ^a	52.80 ^a	1.12	0.41
Lungs (g)	5.88 ^c	9.60 ^b	11.77 ^b	15.03 ^a	16.03 ^a	16.50 ^a	0.22	0.54
Heart (g)	9.34 ^c	11.22 ^{ab}	12.30 ^a	12.50 ^a	13.40 ^a	13.75 ^a	0.61	0.03
Thigh (g)	307.8 ^c	501.0 ^b	558.5 ^b	611.2 ^a	642.5 ^a	655.3 ^a	9.45	0.48
Breast (g)	485.6 ^d	590.3 ^c	730.5 ^b	768.3 ^b	800.2 ^a	813.6 ^a	6.33	0.02
Back (g)	270.4 ^c	388.5 ^b	456.7 ^{ab}	500.6 ^a	516.7 ^a	530.2 ^a	5.90	0.03
Kidney (g)	0.41	0.45	0.43	0.45	0.44	0.45	0.03	0.01
Spleen (g)	2.00	2.01	2.05	2.00	2.36	2.61	0.10	0.01
Pancreas (g)	4.01	4.02	4.56	4.88	4.96	4.98	0.12	0.42
Bursa (g)	3.99	4.08	4.11	4.12	4.18	4.16	0.19	0.80

Results in cells containing various characters vary markedly ($P < 0.05$); dressed weight (DW), dressed percent (DP), and standard error of mean (SEM) *Prosopis africana* oil (PAOs) was added to the diet in the following treatments: T1: control diet + 0% PAOs; T2: basal diet + 2.0 g/kg Neomycin; T3: baseline diet + 200 mg PAOs; T4: baseline diet + 400 mg PAOs; T5: control diet + 600 mg PAOs; and T6: basal diet + 800 mg PAOs.

3.3 Nutrient retention of broiler chickens fed different levels of *Prosopis africana* oil

Table 4 reveals the effect of treatments on nutrient retention of broiler chicks. Dry matter retention was highest for T6 (85.22 %), T5 (84.16 %), T4 (82.80 %), T3 (80.71 %) and lowest for T1 (69.00 %) ($P < 0.05$). Crude protein retention was higher ($P < 0.05$) for T3 (74.61 %), T4 (75.12 %), T5 (77.10 %) and T6 (78.83 %) than for T1 (60.81 %) and T2 (67.10 %). Fat retention was lower ($P < 0.05$) for T1 and T2 than for other treatments. Crude fibre retention was highest ($P < 0.05$) for T3, T4, T5 and T6 and lowest for T1 (38.40 %). Ash values were highest in T3, T4, T5 and T6 and lowest in T1 and T2 ($P > 0.05$). Nitrogen free extract retention was maximum ($P < 0.05$) for T6 (81.34 %), T5 (80.25 %), T4 (80.11 %) and lowest for T3 (78.50 %), T2 (72.15 %) and T1 (70.22 %) ($P < 0.05$). The increased nutritional retention of birds receiving PAOs is consistent with earlier studies (Hernandez *et al.*, 2004), which found that adding plant extracts to a diet increased the apparent whole-tissue digestibility of nutrients. This improvement may have resulted from the activation of digestive enzymes like saliva or bile acids as well as the action of digestive enzymes by PAOs, which increased the number of microbes in the colon. The physiological and chemical functioning of the gastrointestinal system may be impacted by isoprene derivatives, flavonoids, and other plant metabolites. For instance, it has been noted that saponins increase the trypsin enzyme's activity, allowing for effective protein digestion (Rao, 2016). The initial nutrition metabolism may be linked to the stabilizing effect on the gut bacteria (Jamroz *et al.*, 2003). According

to Santi and Kim (2017); Basharat (2018), essential oils have a laxative, spasmolytic, and anti-flatulent action on the gastrointestinal tract. Increased digestive enzyme release improves digestion, the breakdown of metabolic waste products, as well as the absorption and metabolic conversion of the given feed nutrients (Basharat, 2018).

In comparison to alternative treatments, *Prosopis africana* oil improved the digestion of nutrients in birds, improving the animals' overall health (Oloruntola *et al.*, 2019). This research supports the findings of Chowdhury *et al.* (2018), who found that broilers fed diets supplemented with 200 mg/kg of peppermint essential oil had higher crude protein digestibility. Improved feed conversion ratio is required for birds fed PAOs to have improved digestibility. Better muscle accretion occurs in broiler hens due to greater protein digestion (Christine, 2018). On the other hand, Peng *et al.* (2016) found that oregano oil inclusion in the meal at 800 mg/kg had no significant ($P>0.05$) impact on nutritional absorption. According to Symeon *et al.* (2009), adding mixes of oregano, orange peel, and chicory oil to broiler chicken diets at 0.4% did not improve nutrient digestibility. This variation could be explained by variations in dosage, location, as well as bioactive chemical or phytochemical variation.

Table 4: Effect of feeding *Prosopis africana* oil on the nutrient retention of broiler chicks

Parameters	T1	T2	T3	T4	T5	T6	SEM	<i>P</i> -value
Dry matter (%)	69.00 ^c	70.22 ^b	80.71 ^a	82.80 ^a	84.16 ^a	85.22 ^a	5.67	0.34
Crude protein (%)	60.81 ^c	67.10 ^b	74.61 ^a	75.12 ^a	77.10 ^a	78.83 ^a	4.06	0.23
Crude fibre (%)	38.40 ^b	38.99 ^b	40.81 ^a	41.45 ^a	42.72 ^a	43.00 ^a	2.25	0.14
Ether extract (%)	55.63 ^b	58.10 ^b	63.90 ^a	65.16 ^a	65.88 ^a	66.93 ^a	3.98	0.19
Ash (%)	28.19 ^b	28.84 ^b	31.10 ^a	33.85 ^a	33.90 ^a	33.97 ^a	2.06	0.10
NFE (%)	70.22 ^b	72.15 ^b	78.50 ^b	80.11 ^a	80.25 ^a	81.34 ^a	5.15	0.30

Values in cells with various characters vary markedly ($P<0.05$); NFE stands for nitrogen-free extract, and SEM for standard error of mean. T1: base diet plus 0% *Prosopis africana* oil; T2: base diet plus 2.0g/kg neomycin; T3: base diet plus 200 mg PAOs; T4: base diet plus 400 mg PAOs; T5: base diet plus 600 mg PAOs; T6: base diet plus 800 mg PAOs.

Effect of *Prosopis africana* oil on the oxidative stress of broiler chickens

Effect of *Prosopis africana* oil on the oxidative stress of broiler chickens is presented in Table 5. Activities of malondialdehyde (MLAE), superoxide dismutase (SODE), catalase (CATE), reduced glutathione (GSHE) and glutathione peroxidase (GPE) ranged from 6.60 – 14.33 nmol/mL, 100.2 – 238.5 u/mL, 19.48 – 40.11 nmol/g, 5.05 – 9.74 ng/g and 6.21 – 11.44 ng/g respectively. MLA and GPx values were highest in T1; intermediate in T2 and lowest in T3, T4, T5 and T6 ($P<0.05$) whereas SOD and GSH values were higher in T1 and T2 relative to the other treatments ($P<0.05$). CAT value was higher ($P<0.05$) in T3, T4, T5 and T6 than the rest of the treatments. Free radicals are highly reactive, short-lived molecules that can harm cells, including

deoxyribo nucleic acid and cell membranes (Suman *et al.*, 2020; Alagbe, 2019). They are the byproducts of live cells' typical physiological processes (Suman *et al.*, 2020). In the current study, feeding birds with PAOs resulted in considerably higher levels of hepatic antioxidant enzymes and lower levels of malondialdehyde (MDA) than other treatments ($P < 0.05$). Lin *et al.* (2003) noted that consumption of phytogetic feed additives caused an increase in serum antioxidant enzyme activities and a decrease in MDA content, and they reported the same results. Similarly, plant extracts have been shown to boost antioxidant enzyme levels, prevent lipid peroxidation, and scavenge free radicals by Benzie and Wachtel-Galor (2011). On the other hand, Konjufca *et al.* (1997) found that the addition of phytogetic feed ingredients had no impact on the level of blood glutathione. Oluwafemi *et al.* (2021) claim that *Prosopis africana* essential oil creates a range of antioxidant compounds to halt the oxidation of the venerable substrate. Birds can be protected against free radicals by the minerals and vitamins in PAOs, which also function as antioxidants. Numerous physiological processes, such as growth, reproduction, and immunity, have been shown to be negatively impacted by free radical and lipid peroxidation (Gladine *et al.*, 2007; Alagbe and Grace, 2019).

Table 5: Effect of *Prosopis africana* oil on the oxidative stress of broiler chickens

Parameters	T1	T2	T3	T4	T5	T6	SEM	P-value
MLA (nmol/mL)	14.33 ^a	10.95 ^b	8.67 ^c	7.83 ^c	7.11 ^c	6.60 ^c	0.40	0.25
SOD (u/mL)	100.2 ^b	109.6 ^b	113.0 ^b	121.4 ^b	204.6 ^a	238.5 ^a	3.20	0.001
CAT (nmol/g)	17.48 ^c	22.80 ^b	30.80 ^a	35.60 ^a	35.85 ^a	40.11 ^a	0.55	0.13
GSH (ng/g)	5.06 ^b	5.05 ^b	5.03 ^b	5.11 ^b	9.02 ^a	9.74 ^a	0.25	0.30
GPx (ng/g)	6.21 ^c	7.00 ^c	7.07 ^c	8.56 ^c	10.64 ^b	11.44 ^a	0.32	0.11

Means within rows with different letters are significantly different ($p < 0.05$); MLAE: malondialdehyde; SOD: superoxide dismutase; CATE: catalase; GSHE: reduced glutathione; GPE: glutathione peroxidase; SEM: standard error of mean; T1: basal diet + 0 % *Prosopis africana* oil; T2: basal diet + 2.0g/kg Neomycin; T3: basal diet + 200 mg PAOs; T4: basal diet + 400 mg PAOs; T5: basal diet + 600 mg PAOs; T6: basal diet + 800 mg PAOs

4. Conclusion

Ultimately, the experiment's finding showed that dietary supplementation of *Prosopis africana* essential oil (PAOs) did not have any adverse impacts on broiler chickens, such as bacterial resistance or harmful residual deposits in the system. Supplementing oil up to 800 mg/kg had a positive effect on growth performance, feed conversion ratio, immune system and higher carcass output. In order to reconcile the production of animals and the safety of food, PAOs can also be employed.

Funding: "This research received no external funding"

Conflicts of Interest: "The authors declare no conflict of interest."

References

- Suman, T., Arvind, S and Sagar, C. (2021). Phytochemicals and their use in the control of meat oxidation. International Poultry Magazine, pp.13-15.
- Christine, H. (2018). Phytochemicals – a possible and natural alternative to growth enhancing drugs. International Poultry Magazine, pp. 24-27.
- Delphine, M. (2020). Essential oils and organic acids in broilers. International Poultry Magazine, pp.23-25.
- Shittu, M.D., Adejumo, E.O Alagbe, J.O., Ojebiyi, O.O. (2020). Gut morphometric characteristics and ecological response of broiler starter fed varied levels of protein. Asian Journal of Animal Science, 14(1): 33-39.
- Alagbe, J.O. (2017). Effect of dietary inclusion of *Polyalthia longifolia* compared with antibiotics on the performance, carcass characteristics and haematology of broiler chickens. Scholarly Journal of Agricultural Sciences, 7(3): 68-74.
- Tobias, S. (2010). Producers can benefit from essential oil in pig finishing diets. International Poultry Science Magazine, 2010 pp. 20-21.
- Basharat, S. (2018). The potential of phytochemicals in animal nutrition. International Pig Topics 26 (8): 11-13.
- National Research Council (1994). Nutrient requirement of poultry 9th Rev Edn, Washington D.C. National Academy Press.
- Axel, M. (2018). Potential zinc, microflora modulation and broiler meat quality. International Poultry Magazine, pp. 3-5.
- Alagbe, Olujimi John, Oluchi, C.P Agubosi and Rufus, Adebisi Oluwafemi (2023). Histopathology of broiler chickens fed diet supplemented with *Prosopis africana* (African mesquite) essential oil. *Brazilian Journal of Science*, 2(9): 49-59.
- Alagbe, Olujimi John, Oluchi, C. Precious Agubosi., Rufus, Adebisi Oluwafemi., Taiwo, Oladoye Akande., Adegoke, Emmanuel Adegbite and Emiola, I.A. (2023). Haemato-biochemical indices and intestinal microbial population of broiler chickens fed diet supplemented with *Prosopis africana* (African mesquite) essential oil. *Brazilian Journal of Science*, 2(9): 98-110.
- Alagbe, J.O., Bamigboye, S., Nwosu, G.C., Agbonika, D.A and Kadiri Mercy Cincinsoko. (2023). Characterization of bioactive compounds in *Luffa aegyptiaca* leaf ethanolic extracts using gas chromatography and mass spectrometry (GC-MS). *Drug Discovery*, 2023; 17:e10dd1011.
- Alagbe, J.O., Kadiri, M.C., Oluwafemi, R.A., Agubosi, O.C.P and Anorue, D.N. (2023). Analysis of bioactive compounds in ethanolic extracts of *Xylopia aethiopica* leaves using gas chromatography and mass spectrometry technique. *American Journal of Science on Integration and Human Development*, 1(1): 1-10.
- Alagbe, J.O. (2023). Bioactive compounds in ethanolic extract of *Strychnos innocua* root using gas chromatography and mass spectrometry (GC-MS). *Drug Discovery*, 2023; 17:e4dd1005.
- Marc, R. (2010). Healthy pigs with less use of antibiotics – a nutritional approach in three steps. International Pig Magazine, pp. 2-3.
- Alagbe, J.O and Akintayo-Balogun, O.M. (2020). Effects of dietary supplementation of *Albizia lebbek* seed oil (ALO) on the fatty acid composition of weaner rabbits. *Biochemistry and Biotechnology Research*, 8(2): 29-33.
- Jan Dirk, vander Elis and Ester, V. (2013). Phytochemicals – be one step ahead with plant derived feed additives. International Poultry Magazine, pp. 6-8
- Musa, B., Alagbe, J.O., Adegbite Motunrade Betty, Omokore, E.A. (2020). Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2(2):13-21.
- National Population Commission (2006). Nigerian Population Census Report. National Population Commission, Abuja, 21 – 27.

- Suman, T., Arvind, S and Sagar, C. (2014). Phytochemicals and their use in the control of meat oxidation. *International Pig Magazine*, pp. 1-3.
- Sandra, C. (2018). Phyto-genics: how to improve pig production efficiency with plants. *International Pig Magazine*, pp. 5-6.
- Benzoni, G and Guyonvarch, A. (2012). In vitro and in vivo antimicrobial effect of a parented activated clay. *International Poultry Magazine*, pp. 8-10.
- Luis, M.M. (2012). Essential oils and their digestive properties in pigs. *International Pig Magazine*, pp. 4-6.
- Randolf, N. (2018). Influence of phytogenic feed additives. *International Pig Magazine*, pp. 2-3.
- Ines, R. (2010). Phyto-genics: successful AGP replacement in swine feeds. *International Pig Magazine*, pp. 8-9.
- Maria, S. (2016). Gut health management of young piglets with microalgal extracts. *International Pig Magazine*, pp. 3-4.
- Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Sciences* 3(2): 19-29.
- Gberikon, G.M., Agbulu, C.O and Yaji, M.E. (2015). Nutritional composition of fermented powdered *Prosopis africana* soup condiment with or without inocula. *International Journal of Current Microbiology and Applied Sciences*, 4(2): 166-171.
- Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: performance and nutrient retention. *Discovery* 58(314): 134 -142.
- Oluwafemi, R.A., Agubosi, O.C.P and Alagbe, J.O. (2021). Proximate, minerals, vitamins and amino acid composition of *Prosopis africana* (African mesquite) seed oil. *Asian Journal of Advances in Research* 11(1): 21-27.
- Muritala, Daniel Shittu., Alagbe, J.O., Ojebiyi, O.O., Ojediran, T.K and Rafiu, T.A. (2022). Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyspatha longifolia* leaf extract in place of conventional antibiotics. *Animal Science and Genetics* 18(2): 57-71.
- Alagbe, J.O., Adedeji, M.O., Habiba, Z., Nwosu, Gloria and Wyedia Dabara Comfort (2021). Physico-chemical properties of *Indigofera zollingeriana* seed oil. *Asian Journal of Advances in Medical Science* 3(4): 306-308.
- Agubosi, O.C.P., Oluwafemi, R.A and Alagbe, J.O. (2021). The effect of processing on the proximate, mineral and vitamin composition of Neem leaves (*Azadirachta indica*) grown in Gwagwalada, FCT, Abuja. *Abuja Journal of Agriculture and Environment*, 1(1): 293-299.
- Alagbe, J.O (2020). Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and roots of *Indigofera tinctoria*. *International Journal on Integrated Education*. 3(10): 150-157.
- Alagbe, J.O., Shittu, M.D and Eunice Abidemi Ojo (2020). Prospect of leaf extracts on the performance and blood profile of monogastric – A review. *International Journal of Integrated Education*. 3(7): 122-127.
- Adewale, A.O., Alagbe, J.O., Adeoye, Adekemi. O. (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogenic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 3(3): 1-12.
- Singh, A.S., Alagbe, J.O., Sharma, S., Oluwafemi, R.A and Agubosi, O.C.P. (2021). Effect of dietary supplementation of melon (*Citrullus linatus*) seed oil on the growth performance and antioxidant status of growing rabbits. *Journal of Multidimensional Research and Reviews*, 2(1): 78-95.
- Alagbe, J.O (2022). *Prosopis africana* oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Sciences*, 3(2): 19-29.
- Alagbe, J.O., Shittu, M.D and Ushie, F.T. (2021). GC-MS analysis of methanolic stem bark extract of *Zollingeriana indigofera*. *Asian Journal of Advances in Research*, 11(4): 144-146.
- Alagbe, J.O. (2020). Caecal microbial population of growing grasscutters fed *Phyllanthus amarus* and *Piliostigma thonningii* leaf meal mixture as partial replacement for soyameal. *Concept of Dairy and Veterinary Sciences*, 3(5): 350-355.

- Al-Hijazeen, M. (2021). The combination effect of adding rosemary extract and oregano oil on ground chicken meat quality. *Food Science and Technology*, 40: 223-231.
- Agubosi, O.C.P., Oluwafemi, R.A., and Alagbe, J.O. (2021). Preliminary study on GC-MS analysis of *Prosopis africana* seed (African mesquite) oil. *Journal of Ethics and Diversity in International Communication* 1(4): 18-20.
- Lin, X., Jiang, S., Jiang, Z., Zheng, C and Gou, Z. (2003). Effects of equol on H₂O₂-induced oxidative stress in primary chicken intestinal epithelial cells. *Poultry Science* 95:1380–86.
- Jamaroz, M.N.; Sheth, N.R.; Bhatt, V.D. (2003). In Vitro Antibacterial Evaluation of Terminalia chebula as an Alternative of Antibiotics against Bovine Subclinical Mastitis. *Animal Biotechnology*, 30: 151–158.
- Soltan, M.A., Shewita, R.S and El – Katcha, M.I. (2008). Effects of dietary anise seeds supplementation on growth performance, immune response, carcass traits and some blood parameters of broiler chickens. *International Journal of Poultry Science*, 7: 1078–1088
- Schone, F., Vetter, A., Hartung, H., Bergmann, H., Biertümpfel, A., Richter, G., Müller, S and Breitschuh, G. (2006). Effects of essential oils from fennel (*Foeniculi aetheroleum*) and caraway (*Carvi aetheroleum*) in pigs. *Journal of Animal Physiology*, 90: 500–510.
- Oluwafemi, R.A., Uankhoba, I.P and Alagbe, J.O. (2021). Effects of turmeric oil as a dietary supplements on the growth performance and carcass characteristics of broiler chicken. *International Journal of Orange Technologies*, 3(4): 1-9.
- Oluwafemi, R.A., Uankhoba, I.P and Alagbe, J.O. (2021). Effects of turmeric oil as a dietary supplement on the haematology and serum biochemical indices of broiler chickens. *Bioinformatics and Proteomics Open Access Journal* 5(1): 000138
- Hafeez, A., S.M. Suhail, F.R. Durrani, D. Jan, I. Ahmad and A. Rehman (2016). Effect of different types of locally available litter materials on the performance of broiler chicks. *Sarhad J. Agric.* 25(4): 581-586.
- Bamgbose AM, Ogungbenro SD, Obasohan EE, Aruna MB, Oteku IT, Igene UF, Otoikhian CSO, Imasuer JA (2004). Replacement of maize offal/cashew nut for maize in broiler diet. *Proc. 29th Ann Conf. of Nig. Society of Anim. Prod.* 29:219-221.
- Krishan, G and Narang, A. (2014). Use of essential oils in poultry nutrition: A new approach. *Journal of Advanced Veterinary Animal Research*, 1(4): 156-162.
- Hernandez, F., Tsafack Boris Ncedem, Kana Jean Raphaël, Yemdjie Mane Divine, Ebile Dayan Agwah, Ngouana Tadjong Ruben, Donfack Mikael, Tchouan Deffo Gilchrist, Kengni Noubissie Josiane, Teguaia Alexis. (2004) Influence of Two Plant Extracts on Broilers Performance, Digestibility, and Digestive Organ Size. *Poultry Science*, 83, 169-174. <https://doi.org/10.1093/ps/83.2.169>
- Van Veen, T. W. S. (2005). International trade and food safety in developing countries. *Food Control* 16:491–496
- Gladine, C., Morand, C., Rock, E., Bauchart, D., & Durand, D. (2007). Plant extracts rich in polyphenols (PERP). are efficient antioxidants to prevent lipoperoxidation in plasma lipids from animals fed n-3PUFA supplemented diets. *Anim. Feed Sci. Tech.* 136, 281-296.
- Wachtel-Galor, S., Benzie, I.F.F. (2011). Herbal Medicine: an Introduction to its History, Usage, Regulation, Current Trends, and Research Needs. In: second ed. *Herbal Medicine: Biomolecular and Clinical Aspects*. CRC Press/Taylor & Francis, Boca Raton (FL).
- Afub (2021). Effects of phyto-genic feed additives on bird health. *International Poultry Magazine*, 9(8): 13-15.
- Rougiere N and Carre B (2010) Comparison of gastrointestinal transit times between chickens from D+ and D_ genetic lines selected for different digestion efficacy. *Journal Animal Science*. 4: 1861-1872.

- Saeed M., Abbas G., Alagawany M., Kamboh A.A., Abd El-Hack M.E., Khafaga A.F., Chao S. (2019). Heat stress management in poultry farms: a comprehensive overview. *J. Therm. Biology* 84:414–425.
- Alagbe, J.O (2019). Proximate, mineral and phytochemical analysis of *Piliostigma thonningii* stems bark and roots. *International Journal of Biological, Physical and Chemical Studies*, 1(1): 1-7.
- Alagbe, J.O. (2019). Growth performance and haemato-biochemical parameters of broilers fed different levels of *Parkia biglobosa* leaf extracts. *Academic Journal of Life Sciences*. 5(12): 107 – 115.
- Alagbe, J.O., Sadiq, M.R., Anaso, E.U and Grace, F.R. (2019). Efficacy of *Albizia lebbeck* seed oil on the growth performance and carcass characteristics of weaner rabbits. *Sumerianz Journal of Agriculture and Veterinary*. 2(12): 116-122.
- Alagbe, J.O and Oluwafemi, R.A. (2019). Hematology and serum biochemical indices of growing rabbits fed diet supplemented with different levels of *Indigofera zollingeriana* leaf meal. *Progress in Chemical and Biochemical Research*. 2(4): 170-177.
- Alagbe, J.O and Grace, F.R. (2019). Effect of *Albizia lebbeck* seed oil dietary supplementation on the haematological and serum biochemical parameters of weaner rabbits. *Sumerianz Journal of Agriculture and Veterinary*. 2(10): 96 -100.
- Oluwafemi, R.A., Uankhoba, I.P and Alagbe, J.O. (2021). Effects of turmeric oil as adietary supplements on the growth performance and carcass characteristics of broilerchicken. *International Journal of Orange Technologies*, 3(4): 1-9. 35.
- Oluwafemi, R.A., Uankhoba, I.P and Alagbe, J.O. (2021). Effects of turmeric oil as adietary supplement on the haematology and serum biochemical indices of broiler chickens. *Bioinformatics and Proteomics Open Access Journal* 5(1): 000138.36.
- Oluwafemi, R.A., Daniel, S.E and Alagbe, J.O. (2021). Haematology and serumbiochemical indices of broiler chicks fed different inclusion levels of ginger (*Zingiberofficinale*) and garlic (*Allium sativum*) oil mixture. *International Journal of Discoveries and Innovations in Applied Sciences* 1(4): 20-26.37.
- Oluwafemi, R.A., Abdullahi, H and Alagbe, J.O. (2021). Effect of dietary inclusionof ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil mixture on the growthperformance and caecal microbial population of broiler chickens. *Bioequivalence and Bioavailability International Journal* 5(2): 000153.38.
- Alagbe, J.O (2020). Chemical evaluation of proximate, vitamin and amino acidprofile of leaf, stem bark and roots of *Indigofera tinctoria*. *International Journal onIntegrated Education*. 3(10): 150-157.39.
- Alagbe, J.O., Ajagbe, A.D., Attama Jeremiah, Philemon, K.C and Bello, Kamoru, A. (2020). *Albizia lebbeck* stem bark aqueous extract as alternative to antibiotic feedadditives in broiler chick’s diets: Haematology, Serum indices and oxidative status. *International Journal of Biological, Physical and Chemical Studies*, 2(1): 8-15.40.
- Alagbe, J.O (2020). Caecal Microbial Population of Growing Grass Cutters (*Thyronoymys Swinderianus*) Fed *Phyllanthus Amarus* and *Pilogstigma Thonngii* LeafMeal Mixture as Partial Replacement for Soya Bean Meal. *Concept of Dairy andVeterinary Sciences*. 3(5): 350 – 355.41.
- Oluwafemi, R.A., Oluwayinka, E.O and Alagbe, J.O. (2020). Effect of dietarysupplementation of neem oil (*Azadirachtia indica*) on the growth performance andnutrient digestibility of weaned rabbits. *European Journal of Biotechnology and Bioscience*. 8(5): 6-10